

Possible reduction of the total uncertainty on the m_W measurement at LEP2 ^a

J. D'HONDT ^b

Vrije Universiteit Brussel, IIHE, Pleinlaan 2, 1050 Brussel, Belgium

An alternative W mass estimator in $e^+e^- \rightarrow W^+W^- \rightarrow q\bar{q}'\bar{q}q'$ events at LEP2 is designed to optimize the balance between the statistical uncertainty and the systematic uncertainty due to a possible Colour Reconnection effect. The preliminary result for the total uncertainty on the W mass in this channel is roughly 30% lower than those obtained with the standard estimators, based on the SK-I implementation of Colour Reconnection. Also an indirect measurement of the SK-I Colour Reconnection model parameter κ is inferred from the difference between both W mass estimators.

1 Status

A measurement of the mass of the W boson has been performed by all LEP Collaborations¹, this by the method of direct reconstruction of the process $e^+e^- \rightarrow W^+W^- \rightarrow q\bar{q}'\bar{q}q'$, the so-called four jet channel. The uncertainty on the LEP combined $m_W(4q)$ value however is dominated by systematical rather than statistical uncertainties. The major component comes from the assumption in the reconstruction of the event that particles from the decay of different W bosons are independent in the non-perturbative fragmentation phase of the process. Physics phenomena like Colour Reconnection (CR), gluon exchange between fragmentation products, could break this assumption. Neither its theoretical knowledge nor its existence is well established in these processes.

2 Proposal

From the SK-I phenomenological model³ which is implemented in the JETSET fragmentation scheme, we observe that mostly low momentum particles and particles in inter-jet regions are affected by CR and hence influencing the measurant, m_W . Therefore we could design an alternative analysis which is neglecting those particles in the reconstruction of the momenta of the four primary partons, hence decreasing the systematical uncertainty on m_W due to CR but meanwhile increasing the statistical uncertainty because we neglect part of the information content of the event.

Within the standard DELPHI m_W estimator² an iterative procedure was used within each pre-defined jet to find a stable direction of a cone excluding some particles in the calculation of the jet momentum. Starting with the direction of the original jet, the jet direction was recalculated only from those particles which have a opening angle smaller than R_{cone} with this original jet. This process was iterated by constructing a second cone (of the same opening angle)

^aTalk prepared for Rencontres de Moriond XXXVII, electroweak session (Forum for Young Researchers).

^bWork supported by IWT-Vlaanderen.




Figure 1: The left figure shows the dependency of the total uncertainty on m_W and its major components as function of the opening angle of the cone, assuming that all other systematics are invariant under change of R_{cone} . The black line indicates the uncertainty if half of the shift is applied and the other half is quoted as systematic uncertainty. The right figure shows the same as function of the p_{cut} value.

around this new jet direction and the jet direction was recalculated again. The iteration was continued until a stable jet direction was found. The obtained jet momenta were rescaled to conserve the invariant mass of the jets.

The results from Monte Carlo studies using the SK-I Colour Reconnection model with fixed parameter $\kappa = 0.66$ predict a decrease of the total LEP2 uncertainty on $m_W(4q)$ when using the appropriate alternative analysis ($R_{cone} \simeq 0.5\text{rad}$) of around 30%, see Figure 1. Those results are compared with a m_W estimator where simply all particles with momentum below p_{cut} are rejected when calculating the jet kinematics.

An indirect measurement of SK-I model parameter κ is possible from the direct measurement of the difference in reconstructed m_W between the CR-sensitive standard and CR-not-sensitive alternative analysis. From Monte Carlo studies we observe that the statistical significance of this measurement is comparable with the one obtained from the direct measurement. Systematics can however become important.

3 Summary

Alternative exclusive m_W estimators can be designed which have a 30% smaller total uncertainty than the standard ones used by the LEP Collaborations. Also information can be inferred about parametrized Colour Reconnection models by studying the difference between the standard and the proposed alternative m_W estimators.

References

1. The LEP Collaborations, the LEP Electroweak Working Group and the SLD Heavy Flavour Group, LEPEWWG/2002-01.
2. DELPHI Collaboration, *Phys. Lett. B* **511**, 159 (2001).
3. T. Sjöstrand and V. Khoze, *Z. Phys. C* **62**, 281-309 (1994).